DUPLICATE ALSO

Forecasting Research

Forecasting Research Division
Scientific Paper No. 31

Potential vorticities in semi-geostrophic theory

by

Ian Roulstone and Michael J. Sewell

30th January 1995

Meteorological Office
London Road
Bracknell
Berkshire
RG12 2SZ
United Kingdom

ORGS UKMO F
National Meteorological Library
FitzRoy Road, Exeter, Devon. EX1 3PB

Forecasting Research Division
Scientific Paper No. 31

Potential vorticities in semi-geostrophic theory

by

Ian Roulstone and Michael J. Sewell

30th January 1995

Potential vorticities in semi-geostrophic theory

By
Ian Roulstone & Michael J. Sewell*
Forecasting Research Division, Meteorological Office,
London Road, Bracknell, Berkshire. RG12 2SZ.

30th January, 1995.

Abstract

We devise a family of kinematically possible motions in shallow water versions of semi-geostrophic theory, with variable Coriolis parameter, for which two distinct measures of potential vorticity take the same value. We describe conditions for which this value is conserved following the actual motion of a particle. We relate the results to the existing literature, in particular for a constant Coriolis parameter.

1 Introduction

Shallow water theory can serve as a simplified basis for the study of so-called semi-geostrophic theories in geophysical fluid dynamics. We examine shallow water theory and two semi-geostrophic models of that type, with particular reference to a definition of potential vorticity associated with each, and allowing for the Coriolis parameter f to be spatially varying wherever possible. We describe a class of kinematically possible motions under which the expressions for two of the potential vorticities are the same. We give hypotheses under which that value is conserved following the particle.

The plan of the paper is as follows. In §2 we recall the shallow water equations, and the associated measure (q in (6)) of potential vorticity attributed to Rossby (1940, equation (9)). We describe a variational principle for these equations, prompted by Salmon's (1983) work. We state the semi-geostrophic approximation to the shallow water equations for constant f, and we recall (in (12)) the measure of potential vorticity, analogous to Hoskins' (1975, §3(*iii*)) definition, which is conserved by motions which satisfy that approximation. In §3 we describe a generalization, for variable f, of the geostrophic momentum transformation. This leads naturally to the third model, which is a pair of pseudo-hamiltonian

*Permanent address: *Mathematics Department, University of Reading, Whiteknights, P.O. Box 220, Reading. RG6 2AX.*

© Crown Copyright 1995.

equations, also with variable f, studied by Salmon (1985, equations (3.22)), and their associated potential vorticity (Q in (20)).

We are then in a position to exhibit, in §4, a class of kinematically possible motions for which $q = Q$. This our main result. In §5 we establish hypotheses which are sufficient to ensure that both q and Q are conserved following the motion of the particle. This applies, in particular, to the semi-geostrophic equations. Specializing to constant f in §6, we make some specific connections with the previous literature, including Salmon's (1988, equation (5.18)) constraints.

We remark that these results, which stem largely from equation (27), were discovered, in the first instance, using the properties of covariant skew-symmetric tensors, often referred to in the mathematical physics literature as forms, but we have rephrased the proofs in the more elementary form described here. (The calculation about the conservation of q on pp. 70-71 of Rossby (1940) can be understood in terms of these quantities.)

2 Governing equations

The motion of a typical particle in shallow water theory can be described by expressing the current cartesian horizontal coordinates

$$x = x(a, b, t), \quad y = y(a, b, t) \tag{1}$$

as functions, on the right, of the particle labels a, b and the time t. Throughout this paper we shall use the same symbol to denote a function and its generic values, as in this example (1). If $t = 0$ is the reference time, the functions in (1) have the properties $x(a, b, 0) = a$, $y(a, b, 0) = b$.

The incompressibility hypothesis requires the current depth h to be a function $h(a, b, t)$ with the property

$$\frac{h(a, b, 0)}{h(a, b, t)} = \frac{\partial(x, y)}{\partial(a, b)}, \tag{2}$$

where the jacobian on the right is that of the mapping (1). The time derivative of (2) following the particle gives the differential equation of continuity. We regard (2) as describing the explicit solution $h(a, b, t)$ of that continuity equation in terms of (1), and of $h(a, b, 0)$. The latter function of a and b only needs to be given. It is not uncommon to

simplify and normalize by choosing $h(a,b,0) = 1$ in (2), and we make this choice forthwith, but we note that this does constrain the free surface to be parallel to the bed at $t = 0$. In compensation one can take the view that events to be studied take place at much later times.

The equations of horizontal momentum balance for flows over a flat bed which is rotating with position dependent Coriolis parameter $f(x,y)$ are

$$\ddot{x} + g\frac{\partial h}{\partial x} - \dot{y}f = 0, \quad \ddot{y} + g\frac{\partial h}{\partial y} + \dot{x}f = 0\,. \tag{3}$$

Here g is a given constant, representing the combined effect of the acceleration due to gravity and a centrifugal component due to the Earth's rotation. Common choices of the Coriolis function are $f =$ a constant or βy, as approximations to $2\Omega \sin\phi$, depending on purpose. Here β and Ω are constants, and ϕ is latitude, which depends on y, so that x is absent. The superposed dot signifies time differentiation following the particle, i.e. partial differentiation with respect to t when a and b are held constant. The inverse

$$a = a(x,y,t), \quad b = b(x,y,t) \tag{4}$$

of the mapping (1) has been used, with $h(a,b,0) = 1$ in (2), to express $h(a,b,t)$ as another function

$$h(x,y,t) = \frac{\partial(a,b)}{\partial(x,y)}, \tag{5}$$

whose derivatives appear in (3). The basic problem is therefore to solve (3) with (5) for (1), which will then deliver $h(a,b,t)$ from (2).

Let

$$q = \frac{1}{h}\left(\frac{\partial \dot{y}}{\partial x} - \frac{\partial \dot{x}}{\partial y} + f\right) \tag{6}$$

denote a measure of potential vorticity. This name was introduced by Rossby (1940). We shall use q for velocity fields $\dot{x}$, $\dot{y}$ which are *not* solutions of (3) in §§4 and 6.

A result of Salmon (1983, equation (2.20)) can be adapted to show under what conditions (3) are the natural conditions of a variational principle. Let $p(x,y)$ and $r(x,y)$ be any two functions which satisfy the partial differential equation

$$\frac{\partial p}{\partial x} + \frac{\partial r}{\partial y} = f(x,y). \tag{7}$$

The motion (1) can be used to define a functional

$$F[x,y] = \int\int_{t_1}^{t_2} \left[\frac{1}{2}(\dot{x}^2+\dot{y}^2) - r\dot{x} + p\dot{y} - \frac{1}{2}gh\right] \mathrm{d}A_0\mathrm{d}t \tag{8}$$

where the integral is over the area A_0 of label space occupied by the fluid particles under consideration, and over an arbitrary time interval $t_1 \le t \le t_2$. The first variation of (8) for small variations $\delta x(a,b,t)$ and $\delta y(a,b,t)$, taking due account of (2) and (7), is

$$\begin{aligned} \delta F \;=\; & \left[\int((\dot{x}-r)\delta x + (\dot{y}+p)\delta y)\mathrm{d}A_0\right]_{t_1}^{t_2} + \int\int_{t_1}^{t_2} \frac{1}{2}gh^2(l\delta x + m\delta y)\mathrm{d}S\mathrm{d}t \\ & - \int\int_{t_1}^{t_2} \left[(\ddot{x} + g\frac{\partial h}{\partial x} - \dot{y}f)\delta x + (\ddot{y} + g\frac{\partial h}{\partial y} + \dot{x}f)\delta y\right] \mathrm{d}A_0\mathrm{d}t \end{aligned} \tag{9}$$

where S, l, m denote the current boundary of the fluid and its outward unit normal components. Thus whenever the first two terms are zero in (9), we see that (3) implies $\delta F = 0$ and stationary F. It is necessary that, for example, either $h = 0$ or $\delta x = \delta y = 0$ on S.

The semi-geostrophic approximation to equations (3), in the case when f is a constant, is the replacement of the true acceleration by the time derivative of another vector

$$u_g = -\frac{g}{f}\frac{\partial h}{\partial y}, \quad v_g = \frac{g}{f}\frac{\partial h}{\partial x} \tag{10}$$

following the particle. The vector (10) is a notional velocity, called the geostrophic velocity. The semi-geostrophic approximation therefore seeks to find motions (1) satisfying

$$\dot{u}_g + g\frac{\partial h}{\partial x} - \dot{y}f = 0, \quad \dot{v}_g + g\frac{\partial h}{\partial y} + \dot{x}f = 0 \tag{11}$$

with (5) and constant f. Associated with these equations, the expression

$$\frac{1}{h}\left[f + \frac{\partial v_g}{\partial x} - \frac{\partial u_g}{\partial y} + \frac{1}{f}\frac{\partial(u_g, v_g)}{\partial(x,y)}\right] \tag{12}$$

is conserved. This is the shallow water version of Hoskins' (1975) potential vorticity. When f is variable, no measure of potential vorticity associated with (11) is known to be conserved.

3 Transformation to momentum coordinates

The definition (10) of geostrophic velocity suggests a transformation of coordinates

$$X = x + \frac{g}{f^2}\frac{\partial h}{\partial x}, \quad Y = y + \frac{g}{f^2}\frac{\partial h}{\partial y} \tag{13}$$

where (5) is used. However, in this definition we regard f as a *given* function $f(x,y)$ of the *spatial* coordinates in the first instance. Equations (13) are a transformation

$$X = X(x,y,t), \;\; Y = Y(x,y,t) \tag{14}$$

which, at each t, has an inverse

$$x = x(X,Y,t), \;\; y = y(X,Y,t) \tag{15}$$

certainly if

$$\frac{\partial(X,Y)}{\partial(x,y)} \neq 0. \tag{16}$$

Then (15) can be used to write $f(x,y) = f(X,Y,t)$, and the latter will *not* be a *given* function if the motion is unknown in advance. The explicit presence of t in this last expression becomes important in Theorem 2 below. There are some reasons from the literature about constant f (e.g. Eliassen 1948, Hoskins 1975), to call X and Y momentum or geostrophic momentum coordinates.

It is known that (13) is a Legendre (i.e. gradient) transformation when f is constant. This is also true for variable f if and only if

$$\frac{\partial(h,f)}{\partial(x,y)} = 0, \tag{17}$$

i.e. only for certain special motions, and evidently not all motions. Such Legendre transformations have been studied systematically by Chynoweth and Sewell (1989, 1991) for constant f.

Salmon (1985, equations (3.22)) studied certain *generalized* semi-geostrophic equations with pseudo-hamiltonian form in X,Y space, namely

$$\dot{X} = -\frac{1}{f}\frac{\partial H}{\partial Y}, \;\; \dot{Y} = \frac{1}{f}\frac{\partial H}{\partial X} \tag{18}$$

where

$$H(X,Y,t) = \frac{1}{2}(u_g^2 + v_g^2) + gh. \tag{19}$$

Associated with these equations is a *generalized* potential vorticity defined by

$$Q = \frac{f}{h}\frac{\partial(X,Y)}{\partial(x,y)}. \tag{20}$$

Salmon's version (1985, equations (3.12)) of (13) has $f(X,Y)$ written in. Thus (16) is implicitly assumed, but the t dependence in $f(X,Y,t)$ is omitted. When f is a constant, (20) is equivalent to (12).

4 Identification of potential vorticities

We are now in a position to establish the main results of this paper, as follows.

Theorem 1 *There is a class of kinematically possible motions, described in the following proof, and permitting f to vary in physical space, for which*

$$q = Q, \text{ and therefore } \dot{q} = \dot{Q}. \tag{21}$$

Proof

We use the variables X, Y defined by the momentum transformation (13) to construct another transformation

$$\tilde{X} = \tilde{X}(X, Y, t), \ \tilde{Y} = \tilde{Y}(X, Y, t) \tag{22}$$

to new coordinates $\tilde{X}, \tilde{Y}$. The functions on the right of (22) are required to be any which satisfy the partial differential equation

$$\frac{\partial(\tilde{X}, \tilde{Y})}{\partial(X, Y)} = f(X, Y, t). \tag{23}$$

This is one equation, from which to determine two functions, so it will have many solutions. For example, in the special case when f is a non-zero constant, (22) may be any canonical transformation (cf. Sewell and Roulstone, 1993), with t absent.

The momentum transformation (14) can be inserted into the transformation (22) to define a pair of functions $\tilde{X}(x, y, t), \tilde{Y}(x, y, t)$. Let $\lambda(x, y, t)$ be any function, and construct the family of velocity fields $\dot{x}(x, y, t)$, $\dot{y}(x, y, t)$ defined by

$$\left.\begin{aligned} \dot{x} &= r + \frac{\partial \lambda}{\partial x} + \frac{1}{2}\left[\tilde{X}\frac{\partial \tilde{Y}}{\partial x} - \tilde{Y}\frac{\partial \tilde{X}}{\partial x}\right], \\ \dot{y} &= -p + \frac{\partial \lambda}{\partial y} + \frac{1}{2}\left[\tilde{X}\frac{\partial \tilde{Y}}{\partial y} - \tilde{Y}\frac{\partial \tilde{X}}{\partial y}\right], \end{aligned}\right\} \tag{24}$$

By differentiation we see that these satisfy

$$\frac{\partial \dot{y}}{\partial x} - \frac{\partial \dot{x}}{\partial y} = -f + \frac{\partial(\tilde{X}, \tilde{Y})}{\partial(x, y)}, \tag{25}$$

using (7). The chain rule for jacobians gives

$$\frac{\partial(\tilde{X}, \tilde{Y})}{\partial(x, y)} = f\frac{\partial(X, Y)}{\partial(x, y)} = hQ \tag{26}$$

using (23) and (20).

By (6), (25), and (26) we obtain $(21)_1$, and differentiating it following the particle gives $(21)_2$.

□

Summarizing thus far, the two potential vorticities (6) and (20) are the same for any velocity field in the family (24). In other words, (24) is a solution of

$$\frac{\partial \dot{y}}{\partial x} - \frac{\partial \dot{x}}{\partial y} = \left(\frac{\partial(X,Y)}{\partial(x,y)} - 1 \right) f. \tag{27}$$

5 Rate of change of potential vorticity

It follows that either both potential vorticities are conserved, or neither is, during the motions (24).

Theorem 2 *Any solution $X = X(t)$, $Y = Y(t)$ of (18), for* any *hamiltonian function $H(X,Y,t)$ (not only (19)) and for variable $f(X,Y,t)$ $(= f(x,y)$ via (14)), has the property*

$$\dot{Q} = \frac{\partial(X,Y)}{\partial(a,b)} \frac{\partial f}{\partial t}. \tag{28}$$

Here $\partial f/\partial t$ is the partial t derivative of $f(X,Y,t)$, and the jacobian is that of the transformation

$$X = X(a,b,t), \;\; Y = Y(a,b,t) \tag{29}$$

obtained by substituting (1) into (14).

Proof

Combining (5) and (20) gives

$$Q = f \frac{\partial(X,Y)}{\partial(a,b)}. \tag{30}$$

Differentiating following the particle gives

$$\dot{Q} = \dot{f} \frac{\partial(X,Y)}{\partial(a,b)} + f \frac{\partial(\dot{X},Y)}{\partial(a,b)} + f \frac{\partial(X,\dot{Y})}{\partial(a,b)}. \tag{31}$$

The chain rule, with the pseudo-Hamilton equations (18), implies

$$\dot{f} = \frac{\partial f}{\partial X}\dot{X} + \frac{\partial f}{\partial Y}\dot{Y} + \frac{\partial f}{\partial t} = \frac{1}{f}\frac{\partial(H,f)}{\partial(X,Y)} + \frac{\partial f}{\partial t} \tag{32}$$

and

$$\frac{\partial(\dot{X},Y)}{\partial(a,b)} + \frac{\partial(X,\dot{Y})}{\partial(a,b)} = \frac{1}{f^2}\frac{\partial(X,Y)}{\partial(a,b)}\frac{\partial(f,H)}{\partial(X,Y)}. \tag{33}$$

The result (28) follows.

□

When f and Q are constants, (30) shows that (29) is canonical.

Theorem 3

$$\dot{Q} = \dot{q} = 0 \tag{34}$$

for any motion (24) which satisfies the dynamical equations (18), and which also has the property that $\partial f/\partial t = 0$, i.e. for which t is absent from f when expressed as $f(x(X,Y,t), y(X,Y,t))$ in terms of X and Y via (15).

The latter requirement is satisfied a fortiori when f is a constant.

Proof

This is a consequence of Theorems 1 and 2.

□

6 Specializations to f-plane semi-geostrophic theory

Schubert and Magnusdottir (1994) prove that, when f is constant, (27) can be rearranged as a zero curl condition which implies the representation

$$\left.\begin{aligned} \dot{x} &= \frac{\partial\chi}{\partial x} + \frac{1}{2}f\left[(X-x)\frac{\partial Y}{\partial x} - (Y-y)\left(\frac{\partial X}{\partial x}+1\right)\right], \\ \dot{y} &= \frac{\partial\chi}{\partial y} + \frac{1}{2}f\left[(X-x)\left(\frac{\partial Y}{\partial y}+1\right) - (Y-y)\frac{\partial X}{\partial y}\right], \end{aligned}\right\} \tag{35}$$

for arbitrary $\chi(x,y,t)$. It can be verified directly that (35) implies (24) when f is constant. It can also be seen that (35) is included within (24) by making the particular choices

$$p = \frac{1}{2}fx, \quad r = \frac{1}{2}fy, \quad \tilde{X} = |f|^{\frac{1}{2}}X, \quad \tilde{Y} = |f|^{\frac{1}{2}}Y, \quad \lambda = \chi - \frac{1}{2}(xY - yX) \tag{36}$$

when f is a constant.

A brief and elementary calculation using (10) and (13) allows (35), when χ = constant, to be rewritten as

$$\left.\begin{aligned} \dot{x} &= u_g + \frac{1}{2f}\left(u_g\frac{\partial}{\partial x} + v_g\frac{\partial}{\partial y}\right)v_g\,, \\ \dot{y} &= v_g - \frac{1}{2f}\left(u_g\frac{\partial}{\partial x} + v_g\frac{\partial}{\partial y}\right)u_g\,, \end{aligned}\right\} \tag{37}$$

which are precisely Salmon's (1988, equation (5.18)) constraints. The latter *define* his Dirac bracket formulation of f-plane semi-geostrophic theory. When the velocities in (37) are substituted into (6) we obtain (12) identically. Therefore $q = Q$. The potential vorticity Q is conserved and consequently so is q, i.e. $\dot{q} = 0$.

We note that when f is a constant, (13) can be written

$$X = \frac{\partial P}{\partial x},\ Y = \frac{\partial P}{\partial y},\ \text{ with }\ P(x,y,t) = \frac{1}{2}(x^2+y^2) + \frac{g}{f^2}h(x,y,t). \tag{38}$$

The transformation (14) is then of Legendre type, having inverse (15) of the form

$$x = \frac{\partial R}{\partial X},\ y = \frac{\partial R}{\partial Y},\ \text{ with }\ R(X,Y,t) = Xx + Yy - P. \tag{39}$$

This duality leads to a Monge-Ampère equation for P, from (20) and (38), namely

$$Q = \frac{f}{h}\begin{vmatrix} \dfrac{\partial^2 P}{\partial x^2} & \dfrac{\partial^2 P}{\partial x \partial y} \\ \dfrac{\partial^2 P}{\partial y \partial x} & \dfrac{\partial^2 P}{\partial y^2} \end{vmatrix}. \tag{40}$$

In practice, although we have identified Q with the Rossby expression (6), one still has to solve (40) (with suitable boundary conditions) for P, given Q, f and h, in order to calculate X and Y from $(38)_{1,2}$.

Acknowledgements

The authors thank Drs M.J.P. Cullen and M.H. Mawson for useful discussions that prompted and encouraged this work.

References

Chynoweth, S. and Sewell, M.J. (1989). Dual variables in semi-geostrophic theory. *Proc. Roy. Soc. Lond.* Ser. A, **424**, 155-186.

CHYNOWETH, S. and SEWELL, M.J. (1991). A concise derivation of the semi-geostrophic equations. *Q. J. Roy. Met. Soc.*, **117**, 1109-1128.

ELIASSEN, A. (1948). The quasi-static equations of motion. *Geofys. Publ.*, **17**, No. 3.

HOSKINS, B.J. (1975). The geostrophic momentum approximation and the semi-geostrophic equations. *J. Atmos. Sci.*, **32**, 233-242.

ROSSBY, C.-G. (1940). Planetary flow patterns in the atmosphere. *Q. J. Roy. Met. Soc.* **66**, Suppl., 68-87.

SALMON, R. (1983). Practical use of Hamilton's principle. *J. Fluid Mech.*, **132**, 431-444.

SALMON, R. (1985). New equations for nearly geostrophic flow. *J. Fluid Mech.*, **153**, 461-477.

SALMON, R. (1988). Semi-geostrophic theory as a Dirac bracket projection. *J. Fluid Mech.*, **196**, 345-358.

SCHUBERT, W.H. and MAGNUSDOTTIR, G. (1994). Vorticity coordinates, transformed primitive equations and a canonical form for balanced models. *J. Atmos. Sci.*, **51**, 3309-3319.

SEWELL, M.J. and ROULSTONE, I. (1993). Anatomy of the canonical transformation. *Phil. Trans. Roy. Soc. Lond.*, Ser. A, **345**, 577-598.

FORECASTING RESEARCH DIVISION SCIENTIFIC PAPERS
(List valid 30/01/95)
This series of Forecasting Research Division (FR) Scientific Papers . will be papers from all three sections of the Forecasting Research Division i.e. Data Assimilation Research (DA),Numerical Modelling Research (NM), and Observations and Satellite Applications (OB) the latter being formerly known as Nowcasting (NS). This series succeeds the series of Short Range Forecasting Research /Met O 11 Scientific Notes.

1. THE UNIFIED FORECAST /CLIMATE MODEL .
   M.J.P. Cullen
   September 1991

2. Preparation for the use of Doppler wind lidar information in meteorological data assimilation systems
   A.C. Lorenc, R.J. Graham, I. Dharssi, B. Macpherson, N.B. Ingleby, R.W. Lunnon
   February 1992

3. Current developments in very short range weather forecasting.
   B.J. Conway
   March 1992

4. DIAGNOSIS OF VISIBILITY IN THE UK MET OFFICE MESOSCALE MODEL
   AND THE USE OF A VISIBILITY ANALYSIS TO CONSTRAIN INITIAL CONDITIONS
   S.P. Ballard, B.J. Wright, B.W. Golding
   April 1992

5. Radiative Properties of Water and Ice Clouds at Wavelengths
   Appropriate to the HIRS Instrument
   A.J. Baran and P.D. Watts
   2nd June 1992

6. Anatomy of the Canonical Transformation
   M.J. Sewell and I. Roulstone
   27 June 1992

7. Hamiltonian Structure of a Solution Strategy for the Semi-Geostrophic Equations
   I. Roulstone and J. Norbury
   29 June 1992

8. Assimilation of Satellite Data in models for energy and water cycle Research
   A.Lorenc
   July 1992

FORECASTING RESEARCH DIVISION SCIENTIFIC PAPERS

9. The use of ERS-1 data in operational meteorology
A.Lorenc, R.S.Bell, S.J.Foreman, M.W.Holt, D.Offiler
C.D.Hall, D.L.Harrison, S.G.Smith
August 1992

10. Bayesian quality control using multivariate normal distributions
N.B. Ingleby and A. Lorenc
July 1992

11. A NEW APPROACH TO SHALLOW FLOW OVER AN OBSTACLE
I General Theory
A.S. Broad, D. Porter and M.J. Sewell
10 August 1992

12. A NEW APPROACH TO SHALLOW FLOW OVER AN OBSTACLE
II Plane Flow over a Monotonic Mountain
A.S. Broad, D. Porter and M.J. Sewell
10 August 1992

13. A Balanced Ocean Model with Outcropping
Paul Cloke and M.J.P.Cullen
August 1992

14. OSCILLATIONS IN THE ATMOSPHERE'S ANGULAR MOMENTUM AND TORQUES ON THE EARTH'S BULGE
M.J. Bell September 1992

15. Preliminary assessment and use of ERS-1 altimeter wave data
S.J.Foreman, M W Holt and S Kelsall
November 1992

16. NON-GAUSSIAN PROBABILITIES IN DATA ASSIMILATION AND QUALITY CONTROL DECISIONS
Andrew C Lorenc
November 1992

17. Precipitation estimation with AVHRR data: a review
G.L. Liberti
February 1993

18. REAL-TIME CORRECTION OF WEATHER RADAR DATA FOR THE EFFECTS OF BRIGHT BAND, RANGE AND OROGRAPHIC GROWTH
M.Kitchen, R Brown and A G Davies
July 1993

19. A Semi-Geostrophic Ocean Model with Outcropping
Paul Cloke and M.J.P. Cullen
August 25, 1993

FORECASTING RESEARCH DIVISION SCIENTIFIC PAPERS

20. Families of Contact Transformations
M.J. Sewell and I. Roulstone
10 December 1993

21. Global Processing Of Satellite Sounding Radiances
In A Numerical Weather Prediction System
A.J. Gadd, B.R Barwell, S.J. Cox,and R.J. Renshaw
January 1994

22. A Hamiltonian Formulation of the Quasi-Hydrostatic Equations.
I .Roulstone and S.J. Brice
5th April 1994

23. THE CORIOLIS FORCE IN GLOBAL ATMOSPHERIC MODELS
I. THE NAVIER-STOKES EQUATION AND THE HYDROSTATIC PRIMITIVE EQUATIONS
A.A. White, R.A. Bromley, and B.J. Hoskins
April 1994

24. THE CORIOLIS FORCE IN GLOBAL ATMOSPHERIC MODELS
II. DYNAMICALLY CONSISTENT QUASI-HYDROSTATIC EQUATIONS
A.A. White,and R.A. Bromley
April 1994

25. The 1D, Vertical,Version of the Proposed Semi-Implicit Integration Scheme for the Unified Model: Formulation and Test Results.
S. Coulter and M.H. Mawson
October 1994

26. Some issues in numerical methods for the next generation of NWP and climate models
M.P.J. Cullen, T.Davies, M.H. Mawson, J.A. James,
S. Coulter and A. Malcolm
October 1994

27. A Comparison of two vertical grid staggerings
M.J.P. Cullen and.J.James
August 1994

28. On the use of radiosonde humidity observations in midlatitude NWP
A.C. Lorenc, D.Barker, R.S.Bell, B.Macpherson and A.J.Maycock

29. NEW SHALLOW FLOWS OVER AN OBSTACLE
A.S. Broad,D. Porter and M.J. Sewell
19 October 1994

FORECASTING RESEARCH DIVISION SCIENTIFIC PAPERS

30. A shallow-water semi-geostrophic model on the sphere
M.H.Mawson
January 1995

31. Potential vorticities in semi-geostrophic theory
Ian Roulstone and Michael J. Sewell
30th January 1995